

MAXIMIZING COOLING RESOURCES IN TARGETED AREAS

G. E. Dahl
 Department of Animal Sciences
 Institute of Food and Agricultural Sciences
gdahl@ufl.edu
 Georgia Dairy Conference
 18 January 2022

UF IFAS
 UNIVERSITY OF FLORIDA

1

OUTLINE

- Effective cooling approaches
- Water use estimates
- Priority for cooling?
 - Which group first?
 - Lactating? Dry? Calves?
- Summary

UF IFAS
 UNIVERSITY OF FLORIDA

2


EFFECTIVE COOLING

- Goal is 38.6 °C for core temperature
- Combination of water soakers and fans most effective
- Acute versus long term responses – will they match?

KSU via Collier et al., J. Dairy Sci. 89:1244-1253

UF IFAS
 UNIVERSITY OF FLORIDA

3



4


HEAT STRESS DURING LACTATION

- Depresses DMI
- Reduces milk yield
- Recent studies suggest additional metabolic effects beyond DMI
- Recovery dependent on duration, stage of lactation

What about dry cows?

UF IFAS
UNIVERSITY OF FLORIDA

5



6

Gainesville, Florida, USA


- Sand bedded free stalls
- Fans over stalls
- Soakers over feedline
- Fans on at 70° F (21.1°C)
- Soakers on 1 min every 5 min at 72° F



UF IFAS
UNIVERSITY OF FLORIDA

7

STUDY DESIGN: HEAT LOAD OF DRY COWS




UF IFAS
UNIVERSITY OF FLORIDA

De Amaral et al., J. Dairy Sci. 94:86-96

8


HEAT STRESS INCREASES MEAN RECTAL TEMPERATURE




UF IFAS
UNIVERSITY OF FLORIDA

De Amaral et al., J. Dairy Sci. 94:86-96

9



10



11

DRY IN COOL MONTHS IMPROVES PERFORMANCE

Table 1. Milk production and occurrence of mastitis, digestive and respiratory problems, retained fetal membranes, and metritis in cows dried during HOT months (Jun, Jul, Aug) or COOL months (Dec, Jan, Feb) in the first 80 DIM of the subsequent lactation

Item	Dry during HOT months (Jun, Jul, Aug), n = 1,569			Dry during COOL months (Dec, Jan, Feb), n = 1,044			P-value
	Value	Disease ¹	n	Value	Disease ¹	n	
Milk production (kg)	10,351 ± 59.8			10,902 ± 73.3			
Mastitis	0	1,286	82.0	0	950	91.0	0.01
	1	283	18.0	1	94	9.0	
Digestive	0	1,516	96.6	0	973	93.2	0.01
	1	83	5.4	1	71	6.8	
Respiratory	0	1,346	85.9	0	942	90.2	0.01
	1	223	14.2	1	102	9.8	
Retained fetal membranes	0	1,500	95.6	0	1,013	97.0	0.08
	1	89	5.4	1	31	3.0	
Metritis	0	1,500	95.6	0	1,007	96.4	0.35
	1	87	4.2	1	38	3.5	

¹Disease: 0 = cows without the disease; 1 = cows with the disease.

Thompson & Dahl, Prof. Anim. Sci. 28:628-631

UF IFAS
UNIVERSITY OF FLORIDA

12

DRY IN COOL MONTHS IMPROVES REPRODUCTIVE PERFORMANCE

Table 3. Milk production and reproductive performance of cows dried during HOT months (Jun, Jul, Aug) or COOL months (Dec, Jan, Feb) in the first 150 DIM of the subsequent lactation on a commercial farm in Florida

Item	Dry during HOT months (Jun, Jul, Aug)	Dry during COOL months (Dec, Jan, Feb)	P-value
Milk production (kg)	10,547 ± 67.0	11,005 ± 83.38	0.01
Number of breedings (n)	1,048	676	0.03
Mean (no.)	1.59 ± 0.02	1.51 ± 0.03	
DIM to breeding (n)	1,047	676	0.01
Mean (d)	97.0 ± 0.74	91.8 ± 0.92	
DIM to pregnancy (n)	1,051	679	0.01
Mean (d)	131.1 ± 0.85	125.9 ± 1.06	

Thompson & Dahl, Prof. Anim. Sci. 28:628-631

UF IFAS
UNIVERSITY OF FLORIDA

13


LATE GESTATION COOLING

- Do I have to cool cows the entire dry period?
- Do heifers need to be cooled pre-partum?

UF IFAS
UNIVERSITY OF FLORIDA

14


HEAT STRESS INCREASES RECTAL TEMPERATURE AND RESPIRATION RATE




Fabris et al., J. Dairy Sci. 102:5647-5656.

UF IFAS
UNIVERSITY OF FLORIDA


15




16




17




18




19




20




21




22




23



24



25



26

Experiment 2 – No Effect of Colostrum from Cooled or Heat Stressed Cows on Calf Performance

Growth performance of calves born to cows under thermoneutral conditions during the dry period and fed frozen colostrum from cows exposed to either heat stress or cooling during the dry period


Parameter	Heat Stress LSM ± SE	Cooling LSM ± SE	P-value
Birth Weight (kg)	38.8 ± 1.4	39.2 ± 1.5	0.8
Weaning Weight (kg) ¹	68.4 ± 2.5	64.8 ± 2.6	0.4
Prewaning BW Gain (kg) ²	29.6 ± 2.3	25.6 ± 2.4	0.3
Avg. Daily Gain (kg/d)	0.49 ± 0.7	0.43 ± 0.8	0.2
Weaning Withers Height (cm) ¹	84.3 ± 0.8	83.0 ± 0.9	0.4
Prewaning Height Increase (cm) ²	7.8 ± 1.1	6.2 ± 1.0	0.3

¹Weaning weight and weaning height were measured at d 60 of age.
²Prewaning BW gain and height increase was calculated by individually subtracting data at d 60 of age by data at birth.

Monteiro et al., J. Dairy Sci. 97:6426-6439

UF IFAS UNIVERSITY OF FLORIDA

27



28

Heat stress abatement during the dry period: Does cooling improve transition into lactation?
B. C. do Amaral, E. E. Conner, S. Tan, J. Hayes, J. Bullock, and G. E. Dahl
Department of Animal Sciences, University of Florida, Gainesville, 32611
Florida Functional Genomics Laboratory, USDA-ARS, National Agriculture Research Center, Beltsville, MD 20705

Heat stress abatement during the dry period influences metabolic gene expression and improves immune status in the transition period of dairy cows
B. C. do Amaral, E. E. Conner, S. Tan, W. J. Hayes, J. W. Bullock, and G. E. Dahl
Department of Animal Sciences, University of Florida, Gainesville, 32611; National Agriculture Research Center, Beltsville, MD 20705

Effect of heat stress during the dry period on mammary gland development
S. Tan, J. W. Bullock, B. C. do Amaral, L. M. Thompson, W. J. Hayes, S. E. Johnson, and G. E. Dahl
Department of Animal Sciences, University of Florida, Gainesville, 32611

Effect of cooling heat-stressed dairy cows during the dry period on insulin response
S. Tan, L. M. Thompson, A. P. A. Montelero, M. J. Hayes, L. J. Young, and G. E. Dahl
Department of Animal Sciences, University of Florida, Gainesville, 32611; National Agriculture Research Center, Beltsville, MD 20705

Effect of cooling during the dry period on immune response after Streptococcus uberis intramammary infection challenge of dairy cows
L. M. Thompson, S. Tan, A. P. A. Montelero, S. C. Jarvis, and G. E. Dahl
Department of Animal Sciences, University of Florida, Gainesville, 32611

Retrospective analysis of records of calves from 5 studies between 2007 and 2011

Monteiro et al., J. Dairy Sci. 99:8443-8450.

UF IFAS
UNIVERSITY OF FLORIDA

29

Heat stress abatement during the dry period: Does cooling improve transition?
B. C. do Amaral, E. E. Conner, S. Tan, J. Hayes, J. Bullock, and G. E. Dahl
Department of Animal Sciences, University of Florida, Gainesville, 32611
Florida Functional Genomics Laboratory, USDA-ARS, National Agriculture Research Center, Beltsville, MD 20705

Heat stress abatement during the dry period influences metabolic gene expression and improves immune status in the transition period of dairy cows
B. C. do Amaral, E. E. Conner, S. Tan, W. J. Hayes, J. W. Bullock, and G. E. Dahl
Department of Animal Sciences, University of Florida, Gainesville, 32611; National Agriculture Research Center, Beltsville, MD 20705

Effect of heat stress during the dry period on mammary gland development
S. Tan, J. W. Bullock, B. C. do Amaral, L. M. Thompson, W. J. Hayes, S. E. Johnson, and G. E. Dahl
Department of Animal Sciences, University of Florida, Gainesville, 32611

Effect of cooling heat-stressed dairy cows during the dry period on insulin response
S. Tan, L. M. Thompson, A. P. A. Montelero, M. J. Hayes, L. J. Young, and G. E. Dahl
Department of Animal Sciences, University of Florida, Gainesville, 32611; National Agriculture Research Center, Beltsville, MD 20705


Effect of cooling during the dry period on immune response after Streptococcus uberis intramammary infection challenge of dairy cows
L. M. Thompson, S. Tan, A. P. A. Montelero, S. C. Jarvis, and G. E. Dahl
Department of Animal Sciences, University of Florida, Gainesville, 32611

	Bulls	Heifers	Total
Cooling	31	41	72
Heat Stress	30	44	74
Total	61	85	147


Monteiro et al., J. Dairy Sci. 99:8443-8450.

UF IFAS
UNIVERSITY OF FLORIDA

30



31



32

IN UTERO HS DECREASES CALF SURVIVAL

Table 1. Effect of maternal heat stress (HT) or cooling (CL) during late gestation on calf survival

Parameter	CL				HT				P
	AI	IVF	Total	% ^a	AI	IVF	Total	%	
Bull calves, n	30	1	31	---	28	2	30	---	---
Heifer calves, n	29	12	41	---	29	15	44	---	---
DOA ^b	0	0	0.0	---	2	1	3	4.1	0.28
Males mortality by 4 mo of age	1	0	1	3.2	3	0	3	10.0	0.35
Heifers leaving herd before puberty	1	4	5	12.2	3	7	10	22.7	0.26
Due to sickness, malformation or growth retardation	1	0	1	2.4	3	5	8	18.2	0.03
Heifers leaving herd after puberty, before first lactation	1	0	1	2.4	3	0	3	6.8	0.62
Heifers completing first lactation	27	8	35	85.4	22	7	29	65.9	0.05

^aIVF = in vitro fertilization.
^bPercentage of animals (AI + IVF) affected out of total animals (males or females) in the respective treatment.
^cTreatment.
^dDead on arrival. Includes male and female calves.

UF IFAS
UNIVERSITY OF FLORIDA

Monteiro et al., *J. Dairy Sci.* 99:8443-8450.

33

IN UTERO HEAT STRESS DECREASES REPRODUCTIVE PERFORMANCE

Table 2. Effect of maternal heat stress (HT) or cooling (CL) during late gestation on reproductive performance before first lactation of heifers born to HT or CL dams

Parameter	CL	HT	SEM	P
N	36	32	---	---
Age at first AI, mo	13.6	13.8	0.2	0.32
Services per pregnancy d ¹ 30	2.0	2.5	0.2	0.05
Age at pregnancy d ¹ 30, mo	16.1	16.9	0.3	0.07
Services per pregnancy d ¹ 50	2.3	2.6	0.2	0.32
Age at calving, mo	24.8	25.0	0.4	0.72


¹Days after insemination.

Monteiro et al., *J. Dairy Sci.* 99:8443-8450.

UF IFAS
UNIVERSITY OF FLORIDA

34

IN UTERO HEAT STRESS REDUCES MILK PRODUCTION




Monteiro et al., *J. Dairy Sci.* 99:8443-8450.

UF IFAS
UNIVERSITY OF FLORIDA

35


IN UTERO HEAT STRESS DOES NOT AFFECT MATURE BODYWEIGHT




Monteiro et al., *J. Dairy Sci.* 99:8443-8450.

UF IFAS
UNIVERSITY OF FLORIDA


36




37




38




39



40



41



42

TAKE HOME MESSAGES

- Cooling needed for all mature cows – lactating and dry
- Heifers need to be cooled pre-partum to improve yield, protect calf
- Water conservation – esp. “Blue water” - increasingly important consideration for cooling

UF IFAS
UNIVERSITY OF FLORIDA

43
